



Trinational MapBiomias Pampa

Collection 3

Version 1

2023

General Coordinator

Tasso Azevedo

Country coordinators

Diego de Abelleira (Argentina)

Heinrich Hasenack (Brazil)

Santiago Baeza (Uruguay)

Argentina team

Sofía Sarrailhe

Marisa Suklje

Santiago Verón

Santiago Banchemo

Mariano Oyarzabal

Magdalena Bozzola

Brazil team

Eduardo Vélez-Martin

Juliano Schirmbeck

Eliseu José Weber

Uruguay team

Federico Gallego

Maria Vallejos

Andrea Barbieri

Laura Bruzzone

Sebastián Ramos

Gonzalo Rama

1 INTRODUCTION

1.1 Scope and content of the document

The objective of this document is to describe the theoretical basis, justification and methods applied to produce annual maps of land use and land cover (LULC) in the South American Pampa biome of Argentina, Brazil and Uruguay from 1985 to 2022 (Collection 3). The document presents a general description of the satellite image processing, the feature inputs and the process step by step applied to obtain the annual classifications.

1.2 Region of Interest

Trinational Pampa MapBiomass initiative was created to produce LULC annual maps for the Pampa biome corresponding to Argentina, Brazil and Uruguay territories. Other phytogeographic regions closed or interspersed with Pampa were partially added to allow a better regional delimitation. Thus, a neighboring area of *Espinal* phytogeographic province, around the Pampa biome, as well as the Paranaense phytogeographic province, located both in Argentina were also included (**Figure 1**).

The total mapped area was 109,21 million hectares (Mha), being 83,81 Mha in the Pampa (21%), 23,15 Mha in the Espinal (77%) and 2,25 Mha in the Paraná river delta (2%).




Figure 1. Region of interest mapped in the Trinational MapBiomias Pampa initiative (collection 3), including the typical areas of the Pampa, Espinal, and Parana river Delta.

2 GEOGRAPHICAL UNITS OF CLASSIFICATION

In each country, the classification process was carried out in smaller spatial units. These units correspond to subregional homogeneous zones based on several criteria, nationally defined, including geomorphology, soils, vegetation types and land use patterns.

The study area was divided in 27 homogeneous zones, thirteen in Argentina, seven in Brazil and seven in Uruguay (**Figure 2**).

The purpose of these geographical units of classification was to try to reduce samples and classes confusion and to allow a better balance of samples and results to improve accuracy.




Figure 2. Country defined homogeneous subregions used in the classification process of the South American Pampa biome.

3 REMOTE SENSING DATA

3.1 Landsat Collection

The imagery dataset used in the *Trinational Pampa MapBiomias* Collection 3 was obtained from the Landsat sensors Thematic Mapper (TM), Enhanced Thematic Mapper Plus (ETM+) and the Operational Land Imager and Thermal Infrared Sensor (OLI-TIRS), on board of Landsat 5, Landsat 7 and Landsat 8, respectively. The Landsat imagery collections with 30 m-pixel resolution were accessible via Google Earth Engine, and were provided by NASA and USGS. The *Trinational Pampa MapBiomias* Collection 3 used Collection 2, Tier 1 Landsat Surface Reflectance products from USGS, which underwent through radiometric calibration and orthorectification correction based on ground control points and digital elevation model to account for pixel co-registration and correction of displacement errors. A total of 74 scenes were used to cover the entire region, where each of them is totally or partially within the area.

According to the year and the quality of available images, a specific Landsat collection was selected:

- from 1985 to 1999: Landsat 5,
- year 2000: Landsat 5 (Brazil and Uruguay) and Landsat 7 (Argentina),
- years 2001, 2002 and 2012: Landsat 7,
- from 2003 to 2011: Landsat 5,
- from 2013 to 2022: Landsat 8.

3.2 Landsat Mosaics

All Landsat scenes were merged and clipped within standardized spatial units for data processing, hereafter called ‘charts’, based on the grid of the World International Chart to the Millionth, at the 1:250,000 scale level. A total of 99 charts were used to cover the biome (**Figure 3**). Each chart sets the geographical limits to build up the temporal and spatial Landsat mosaics and to proceed with digital classification procedures. Each geographical classification unit was generated by merging the correspondent mosaic charts.




Figure 3. Charts scheme used to build up Landsat mosaics used throughout the classification process.

3.3 Definition of the temporal period

The mosaics were formed by the composition of pixels in each set of images for a certain time period. The periods of the year in which the images are selected vary by country and result from the balance between the probability of maximizing the differences in classes spectral behavior and the availability of cloud-free images. In Uruguay and Brazil, the considered period was from September to November of each year, while in Argentina from May to July. Nevertheless, for some years this period was adapted (extended one to three months) for each chart according to the availability of cloud-free images. For example, if during the three-months period a cloud free mosaic could not be generated, the period was extended to four, five or six months to get a complete or almost complete mosaic.

For the selection of Landsat scenes a threshold of 90% of cloud cover was applied (i.e., any available scene with up to 90% of cloud cover was accepted). This limit was established based on a visual analysis, after many trials observing the results of the cloud removing/masking algorithm. Time periods were extended for some years and portions of the study area when the availability of cloud-free images was low.

4 CLASSIFICATION

4.1 Overview of methodological process

The methodological procedures of Collection 3 included several steps (**Figure 4**). The first step was to generate annual Landsat image mosaics based on yearly periods. The second step was to generate a new selection of temporally stable samples derived from the stable areas of the maps of Collection 2. Stable areas were defined in sub-periods of near 10 years-length (1985-1994, 1995-2004, 2005-2014 and 2015-2022). Then, the spectral feature inputs derived from the Landsat bands were extracted and associated to each sample point. Once the samples for each LULC class were selected for each of the subregions, it was possible to adjust the training data set according to its statistical needs. The number of samples for training for each class was defined initially according to the proportion of the area of each class and its variation along the classification period (sample size balance). Additionally, to improve the classification results, complementary samples were generated, defining georeferenced points of different classes by visual interpretation of historical satellite images (high and very high resolution images) and time series of vegetation indices.

Based on the adjusted training data set, a supervised classification using the random forest algorithm was run.

Following that, gap, spatial, temporal and frequency filters were applied to remove classification noise and stabilize the classification. The LULC maps of each subregion were integrated to generate the final map of Collection 3. The MapBiomas annual LULC maps were used to derive the transition analysis (with an additional spatial filter application) and statistics. The statistical analysis covered different spatial territories, such as countries, state similar and municipality similar levels of each country, water basin and phytogeographic provinces.














Figure 4. Classification process of Collection 3 in the *MapBiomas South American Pampa* biome for the period 1985-2022.

4.2 Map Legend

The classification for the *Trinational Pampa MapBiomas* initiative using Landsat mosaics included nine land use and land cover (LULC) classes (**Table 1**): Forest formation (3), Savanna formation (4), Wetland (11), Grassland (12), Agriculture or pasture (21), Silviculture (9), Non vegetated area (22), River, lake or ocean (33) and

Not observed (27). In Argentina, Agriculture or pasture class (21) was also separated in Agriculture (18) and Pasture (15) and this data was also made available on an alternative map. A full description of the legend is described in the [document Legend Description](#).

Table 1. Land cover and land use classes considered for digital classification of Landsat mosaics for the South American Pampa biome - Collection 3.

Legend class of Collection 3	Numeric ID	Color
1.1. Forest formation	3	
1.2. Savanna formation	4	
2.1. Wetland	11	
2.2. Grassland	12	
3.1. Pasture	15	
3.2. Agriculture	18	
3.3. Agriculture or pasture	21	
3.4. Forest plantation	9	
4. Non vegetated area	22	
5.1. River, lake or ocean	33	
6. Not observed	27	

4.3 Annual Mosaics

The total available bands of the Trinational MapBiomass Pampa feature space is composed of 93 input variables, including the original Landsat bands, fractional and textural information derived from these bands (**Table 2**). Reducers were used to generate temporal features such as:

- Median: median of the pixel values of the best mapping period defined by each country.
- Median_dry: median of the quartile of pixels with the lowest NDVI values.
- Median_wet: median of the quartile of pixels with the highest NDVI values.
- Amplitude: amplitude of variation of the index considering all the images of each year.
- stdDev: standard deviation of all pixel values of all images of each year.
- Min: lower annual value of the pixels of each band.

Table 2. List of the variables included in the feature space used in the classification of the Mapbiomas Trinational Pampa Landsat image mosaics in the Collection 3 (1985-2022).

ID	Variable	Description	Statistics	Temporal range	Script acronym	Group
0	Evi 2	Enhanced Vegetation Index 2	amplitude	mosaic months	evi2_amp	Spectral index
1	Gv	Green vegetation fraction	amplitude	mosaic months	gv_amp	Spectral Mixture Modeling
2	Ndfi	Normalized Difference Fraction Index	amplitude	mosaic months	ndfi_amp	Spectral Mixture Modeling
3	Ndvi	Normalized Difference Vegetation Index	amplitude	mosaic months	ndvi_amp	Spectral index
4	Ndwi	Normalized Difference Water Index	amplitude	mosaic months	ndwi_amp	Water Index
5	Soil	Soil fraction	amplitude	mosaic months	soil_amp	Spectral Mixture Modeling
6	Wefi	Woodland ecosystem fraction index	amplitude	mosaic months	wefi_amp	Fraction index
7	Blue	Landsat band	median	mosaic months	blue_median	Landsat band
8	Blue dry	Landsat band	median	year -first quartile	blue_median_dry	Landsat band
9	Blue wet	Landsat band	median	year – fourth quartile	blue_median_wet	Landsat band
10	Cai	Cellulose Absorption Index	median	mosaic months	cai_median	Spectral index
11	Cai dry	Cellulose Absorption Index	median	year -first quartile	cai_median_dry	Spectral index
12	Cloud	Cloud fraction	median	mosaic months	cloud_median	Spectral Mixture Modeling
13	Evi 2	Enhanced Vegetation Index 2	median	mosaic months	evi2_median	Spectral index
14	Evi 2 dry	Enhanced Vegetation Index 2	median	year -first quartile	evi2_median_dry	Spectral index
15	Evi 2 wet	Enhanced Vegetation Index 2	median	year – fourth quartile	evi2_median_wet	Spectral index
16	Gcvi	(nir/green – 1)	median	mosaic months	gcvi_median	Spectral index
17	Gcvi dry	(nir/green – 1)	median	year -first quartile	gcvi_median_dry	Spectral index
18	Gcvi wet	(nir/green – 1)	median	year – fourth quartile	gcvi_median_wet	Spectral index
19	Green	Landsat band	median	mosaic months	green_median	Landsat band
20	Green dry	Landsat band	median	year -first quartile	green_median_dry	Landsat band
21	Green wet	Landsat band	median	year – fourth quartile	green_median_wet	Landsat band
22	Gv	Green vegetation fraction	median	mosaic months	gv_median	Spectral Mixture Modeling
23	Gvs	GV / (100 - shade)	median	mosaic months	gvs_median	Spectral Mixture Modeling
24	Gvs dry	GV / (100 - shade)	median	year -first quartile	gvs_median_dry	Spectral Mixture Modeling
25	Gvs wet	GV / (100 - shade)	median	year – fourth quartile	gvs_median_wet	Spectral Mixture Modeling
26	Hallcover	Hall cover vegetation index	median	mosaic months	hallcover_median	Spectral index
27	Ndfi	Normalized Difference Fraction Index	median	mosaic months	ndfi_median	Spectral Mixture Modeling
28	Ndfi dry	Normalized Difference Fraction Index	median	year -first quartile	ndfi_median_dry	Spectral Mixture Modeling
29	Ndfi wet	Normalized Difference Fraction Index	median	year – fourth quartile	ndfi_median_wet	Spectral Mixture Modeling
30	Ndvi	Normalized Difference Vegetation Index	median	mosaic months	ndvi_median	Spectral index
31	Ndvi dry	Normalized Difference Vegetation Index	median	year -first quartile	ndvi_median_dry	Spectral index
32	Ndvi wet	Normalized Difference Vegetation Index	median	year – fourth quartile	ndvi_median_wet	Spectral index
33	Ndwi	Normalized Difference Water Index	median	mosaic months	ndwi_median	Water Index

ID	Variable	Description	Statistics	Temporal range	Script acronym	Group
34	Ndwi dry	Normalized Difference Water Index	median	year -first quartile	ndwi_median_dry	Water Index
35	Ndwi wet	Normalized Difference Water Index	median	year – fourth quartile	ndwi_median_wet	Water Index
36	Near Infrared (NIR)	Landsat band	median	mosaic months	nir_median	Landsat band
37	Near Infrared (NIR) dry	Landsat band	median	year -first quartile	nir_median_dry	Landsat band
38	Near Infrared (NIR) wet	Landsat band	median	year – fourth quartile	nir_median_wet	Landsat band
39	Npv	Non-photosynthetic vegetation fraction	median	mosaic months	npv_median	Spectral Mixture Modeling
40	Pri	Photochemical reflectance index	median	mosaic months	pri_median	Spectral index
41	Pri dry	Photochemical reflectance index	median	year -first quartile	pri_median_dry	Spectral index
42	Pri wet	Photochemical reflectance index	median	year – fourth quartile	pri_median_wet	Spectral index
43	Red	Landsat band	median	mosaic months	red_median	Landsat band
44	Red dry	Landsat band	median	year -first quartile	red_median_dry	Landsat band
45	Red wet	Landsat band	median	year – fourth quartile	red_median_wet	Landsat band
46	Savi	Soil-adjusted Vegetation Index	median	mosaic months	savi_median	Spectral index
47	Savi dry	Soil-adjusted Vegetation Index	median	year -first quartile	savi_median_dry	Spectral index
48	Savi wet	Soil-adjusted Vegetation Index	median	year – fourth quartile	savi_median_wet	Spectral index
49	Sefi	Savanna Ecosystem Fraction Index	median	mosaic months	sefi_median	Fraction index
50	Sefi dry	Savanna Ecosystem Fraction Index	median	year -first quartile	sefi_median_dry	Fraction index
51	Shade	Shade fraction	median	mosaic months	shade_median	Spectral Mixture Modeling
52	Soil	Soil fraction	median	mosaic months	soil_median	Spectral Mixture Modeling
53	Shortwave Infrared (SWIR) 1	Landsat band	median	mosaic months	swir1_median	Landsat band
54	Shortwave Infrared (SWIR) 1 dry	Landsat band	median	year -first quartile	swir1_median_dry	Landsat band
55	Shortwave Infrared (SWIR) 1 wet	Landsat band	median	year – fourth quartile	swir1_median_wet	Landsat band
56	Shortwave Infrared (SWIR) 2	Landsat band	median	mosaic months	swir2_median	Landsat band
57	Shortwave Infrared (SWIR) 2 dry	Landsat band	median	year -first quartile	swir2_median_dry	Landsat band
58	Shortwave Infrared (SWIR) 2	Landsat band	median	year – fourth quartile	swir2_median_wet	Landsat band

ID	Variable	Description	Statistics	Temporal range	Script acronym	Group
	wet					
59	Wefi	Woodland ecosystem fraction index	median	mosaic months	wefi_median	Fraction index
60	Wefi wet	Woodland ecosystem fraction index	median	year – fourth quartile	wefi_median_wet	Fraction index
61	Blue min	Landsat band	minimum	mosaic months	blue_min	Landsat band
62	Green min	Landsat band	minimum	mosaic months	green_min	Landsat band
63	Near Infrared (NIR) min	Landsat band	minimum	mosaic months	nir_min	Landsat band
64	Red min	Landsat band	minimum	mosaic months	red_min	Landsat band
65	Shortwave Infrared (SWIR) 1	Landsat band	minimum	mosaic months	swir1_min	Landsat band
66	Shortwave Infrared (SWIR) 2	Landsat band	minimum	mosaic months	swir2_min	Landsat band
67	Blue	Landsat band	standard deviation	mosaic months	blue_stdDev	Landsat band
68	Cai	Cellulose Absorption Index	median	mosaic months	cai_stdDev	Spectral index
69	Cloud	Cloud fraction	standard deviation	mosaic months	cloud_stdDev	Spectral Mixture Modeling
70	Evi 2	Enhanced Vegetation Index 2	standard deviation	mosaic months	evi2_stdDev	Spectral index
71	Gcvi	(nir/green – 1)	standard deviation	mosaic months	gcvi_stdDev	Spectral index
72	Green	Landsat band	standard deviation	mosaic months	green_stdDev	Landsat band
73	Gv	Green vegetation fraction	standard deviation	mosaic months	gv_stdDev	Spectral Mixture Modeling
74	Gvs	GV / (100 - shade)	standard deviation	mosaic months	gvs_stdDev	Spectral Mixture Modeling
75	Hallcover	Hall cover vegetation index)	standard deviation	mosaic months	hallcover_stdDev	Spectral index
76	Ndfi	Normalized Difference Fraction Index	standard deviation	mosaic months	ndfi_stdDev	Spectral Mixture Modeling
77	Ndvi	Normalized Difference Vegetation Index	standard deviation	mosaic months	ndvi_stdDev	Spectral index
78	Ndwi	Normalized Difference Water Index	standard deviation	mosaic months	ndwi_stdDev	Water Index
79	Near Infrared (NIR)	Landsat band	standard deviation	mosaic months	nir_stdDev	Landsat band

ID	Variable	Description	Statistics	Temporal range	Script acronym	Group
80	Red	Landsat band	standard deviation	mosaic months	red_stdDev	Landsat band
81	Savi	Soil-adjusted Vegetation Index	standard deviation	mosaic months	savi_stdDev	Spectral index
82	Sefi	Savanna Ecosystem Fraction Index	standard deviation	mosaic months	sefi_stdDev	Fraction index
83	Shade	Shade fraction	standard deviation	mosaic months	shade_stdDev	Spectral Mixture Modeling
84	Soil	soil fraction	standard deviation	mosaic months	soil_stdDev	Spectral Mixture Modeling
85	Shortwave Infrared (SWIR) 1	Landsat band	standard deviation	mosaic months	swir1_stdDev	Landsat band
86	Shortwave Infrared (SWIR) 2	Landsat band	standard deviation	mosaic months	swir2_stdDev	Landsat band
87	Wefi	Woodland ecosystem fraction index	standard deviation	mosaic months	wefi_stdDev	Fraction index
88	Slope	Terrain slope	identity	Permanent	slope	Geomorphometric
89	Green Texture	Texture from Landsat band	mean	mosaic months	green_median_texture	
90	Latitude	Geographical coordinate	-	Permanent	Latitude	Geographic
91	Longitude	Geographical coordinate	-	Permanent	Longitude	Geographic
92	Ndvi_3years	Normalized Difference Vegetation Index	amplitude	Last 3 years mosaic months	ndvi_amp_3y	Spectral index

4.4 Classification algorithm, training samples and parameters

Classification was performed subregion by subregion, year by year, using the Random Forest algorithm (Breiman, 2001) available in Google Earth Engine, running 100 iterations (random forest trees).

Training samples for each subregion were defined following a strategy of using random pixels for which the land use and land cover remained the same (stable samples) along the maps of Collection 2 over different subperiods: 1985-1994, 1995-2004, 2005-2014 and 2015-2022, named as “stable samples”.

The identification of stable areas to extract random pixels or “stable samples” was based on a criterion of minimum temporal frequency aiming to ensure confidence to use them as training areas. Each pixel should be classified with the same LULC class for at least a minimum number of years within each sampling subperiod (1985-1994, 1995-2004, 2005-2014 and 2015-2022). The thresholds for some classes and each country and subregion were not the same. A layer of pixels with a stable classification for each subperiod was then generated by applying such thresholds. From the resulting layer of stable samples, a subset of 2,000 samples for each subregion were randomly generated for each class for each year. It is important to clarify that not all of these samples were necessarily used in the classification process.

In addition, a classical procedure to detect outliers was implemented. For each year, and within each training class, we searched for outliers in all variables. An outlier was defined as any value of a specific variable lower or higher than 1.5 times the interquartile range (the first quartile value subtracted from the third quartile value) considering all values of this variable within a specific class of a particular year. Samples containing values considered outliers for some variables were not discarded a priori, but fixed by replacing those values with the 5th percentile or the 95th percentile, whenever they were lower or higher than the thresholds considered, respectively. Finally, we disregarded only those samples containing simultaneously more than 6 variables of the feature space with values considered as outliers.

4.4.1 Sample size balance

We generated a fixed number of samples for each class and subregion of classification. However we used in the classification process only a random subset based on the class area proportion within each subregion, considering each year to

be classified. To do this we previously adjusted linear simple functions to estimate the area of each class for each year from 1985 to 2022, based on the annual class area observed along the Collection 2 dataset. These functions were used to estimate, for each year, the proportion of each class to train the classifier. Then, these annual proportions for each class were set to extract a subset of the available samples for the correspondent classification in each year. Whenever the classification resulted in overestimation or underestimation of the class after comparing with supplemental information (e.g.: Collection 2 maps, independent crop type maps, etc.) this proportion was adjusted changing the bias (intercept of linear regression model) accordingly. Notwithstanding the above, a minimum number of 50 to 100 samples per class was set for each region and year, to ensure the correct detection of the less frequent categories.

4.4.2 Complementary samples

The need for adding complementary samples was evaluated by visual inspection of the output of a preliminary classification, with both Landsat and high-resolution images available in GEE and time series of vegetation indices, and also by comparing with the Collection 2 classification. Complementary sample collection was also done manually using points in Google Earth Engine Code Editor. All the false-color images of the 38 years (1985-2022) Landsat mosaics and the vegetation index time series were checked at the selected point. Based on the knowledge of each subregion, the samples for different classes were collected.

4.4.5 Final classification

The final classification was performed for all subregions and years combining stable and complementary samples. For some years the classification output resulted in anomalous results for some classes. Then, it was necessary to improve the classification through a new sample size balance and a specific set of complementary samples.

4.4.6 Post-classification

The results of the final classification were improved through a sequence of filters, to correct missing data, “salt-and-pepper” classification errors and, specially, cases of misclassification. Temporal filters were done with the aim to generate a more stable classification pattern over time, avoiding unexpected classes variation during short times.

4.4.6.1. Gap fill filter

A filter to fill no-data pixels (“gaps”) was applied. Because theoretically the no-data values are not allowed, they are replaced by the temporally nearest valid classification. In this procedure, if no “future” valid position was available, then the no-data value was replaced by its previous valid class. Therefore, gaps should only exist if a given pixel has been permanently classified as no-data throughout the entire temporal domain.

4.4.6.2. Spatial filter

The spatial filter avoids unwanted modifications to the edges of the pixel groups, a spatial filter was built based on the "connectedPixelCount" function. Native to the GEE platform, this function locates connected components (neighbors) that share the same pixel value. Thus, only pixels that did not share connections to a predefined number of identical neighbors were considered isolated. In this filter, at least six connected pixels were needed to reach the minimum connection value. Consequently, the minimum mapping unit is directly affected by the spatial filter applied, and it was defined as 6 pixels (~0,5 ha).

4.4.6.3. Temporal filters

The temporal filters use the information from the year before and after to identify and correct a pixel misclassification, considered as cases of invalid transitions. In a first step, the filter looks for specific cover classes (3, 4, 11, 12, 33) that are not this class in 1985 and were kept unchanged in 1986 and 1987 and then corrects the 1985's value to avoid any regeneration in the first year. In a second step, the filter looks at a pixel value in 2022 that for example is not 21 (Agriculture or pasture) but is equal to 21 in 2020 and 2021. The value in 2022 is then converted to 21 to avoid any regeneration in the last year. The third process looks in a 3-year moving window to correct any value that changed in the middle year and returns to the same class next year.

A temporal filter with a slightly different approach was applied to solve problems in forestry classification. To correct the problems related to the years with forestation cutting, interrupting a continuous series of years classified as forestry we used a special six-year spatial filter. The rule of application checks whether two years before and two years after the class was forestation, if this is true it shifts the classification of the two middle years to silviculture.

4.4.6.4. Frequency filter

To correct classification problems associated with some classes in specific regions, frequency filters were applied to use the temporal information available for each pixel to correct cases of false positives. The general logic of the frequency filter is to search for each pixel a specific combination of classes throughout the 38 years producing a subset of pixels considered eligible for correction. Then the filter detects and overwrites only those years where cases of false positives are present using a fixed class value, that usually is the mode of classifications detected along the temporal range. This type of filter were used with parsimony to solve very well delimited cases.

4.4.6.5. Specific filters

Additional specific filters were generated to remove unexpected classification changes that remained after applying previous standard filters. In general, these filters that we applied work with frequency and incidence. Frequency is the number of years a class occurs in a pixel. The incidence is the number of times that a pixel classification changes along the entire series of years. The application of these filters was limited to fix problems of false transitions between specific classes.

We also used a filter that eliminates problems related to the shadows of the mountains. These filters use characteristics of the relief, in addition to the frequency to be applied. It corrects false positives of water and wetland in shaded slopes in regions with wavy relief. The filter selects all pixels classified as water at least in one year but in less than 34 years (<95%), or as wetland at least in one year but in less than 32 years (<90%), whenever occurring in areas of cliffs and slopes, established by a combination of slope data (SRTM derived) with HAND (Height Above the Nearest Drainage) database, to define places where it is not expected the presence of water or wetland. In such cases, both classes were replaced by the class corresponding to the pixel mode.

A filter to smooth abrupt transitions between the first and the second year (198-1986) and the last and penultimate years (2021-2022) was applied only in Argentina and Brazil. It has been observed in previous collections, that the last year of the series registered an unexpected increase in the area of anthropic classes and a decrease of natural classes, most likely corresponding to an artifact resulting from the set of applied filters. To alleviate the problem, a filter was developed to smooth this abrupt transition, avoiding all transitions from natural areas to anthropic areas, and vice versa, in

patches equal to or smaller than 2 hectares. In these cases, the correspondent pixels from the last year receive the same classification as the penultimate year as well as pixels from the first year receive the same classification as the second year.

Exceptionally, the spatial effect of some filters was limited set of polygons, in such a way as not to modify the entire zone classification. Examples of these filters include: a grassland filter that unifies wet and dry years, taking into account the coverage of that place and not the rainfall of a particular year. Or a rice filter that corrects sites classified as wet grasslands, only for certain years, as long as it has been previously classified as agriculture.

5 VALIDATION STRATEGIES

Validation was performed for the classifications of the years 1986, 2001 and 2018 following the good practices recommendations proposed by Olofsson et al. (2014) for area and error estimation. A total of 2,330 samples were defined as necessary for the analysis. The number of samples for each class was proportional to the area of each class obtained from Collection 1 for the year 2010. Independent samples were raffled and class classified by visual interpretation of Landsat images, very high resolution images from Google Earth and time series of vegetation indices. Two interpreters evaluated each of the sample points generated from the stratified random design. In those sample points where discordance in class classification was detected among interpreters, a third interpreter defined the final class assignment. More details of the validation methodology are described in Baeza et al. (2022). When a final class could not be defined by the three interpreters (e.g. three different class assignments), a final class was agreed by a team of interpreters.

Validation results showed an overall accuracy of 75% for 1986, 79 % for 2001 and 80% for 2018. In all three cases, most of the associated errors were location mismatches rather than quantity mismatches (see Pontius and Milloes, 2011), allowing for more precise area estimates (global accuracy + location mismatch): 91%, 93% and 90% for the years 1986, 2001 and 2018, respectively. **Tables 3, 4 and 5** show the contingency matrices performed to evaluate the outcome of the classifications of collection 3 for the years 1986, 2001 and 2018 from the independent data set explained above. **Figure 5** shows the user and producers accuracies (the opposite of

commission and omission errors, respectively) of each category. The lowest accuracies (in percentage) occurred in the forestation class in 1986, mainly due to confusion with natural woody cover. This was associated with the low proportion of forest plantations in that year and the difficulty of discriminating between these coverages without higher resolution images. Another important confusion was associated with the mixing of wetlands with grasslands, associated with the change in flooding levels of floodable grasslands. The largest number of points incorrectly classified (by inclusion or omission) was due to confusion between grassland and the agriculture/pasture class. The high physiognomic similarity and the intra class heterogeneity of the spectral response of both natural grasslands (different communities, landscape positions, location in the study area) and sown pastures (different species, sowing dates, pasture age, etc.) generate an overlapping of the spectral signatures of both coverages, which explains the confusion in some of the evaluated points.

Table 3. Contingency matrix resulting from comparing the 1986 map with the independent data set of the same year.

		Reference							Total
		Classes	Natural woody vegetation	Forest plantation	Wetland	Grassland	Agriculture or pasture	Non-vegetated area	
Classification	Natural woody vegetation	145	7	5	24	6	0	0	187
	Forest plantation	3	6	1	2	1	0	0	13
	Wetland	8	1	106	6	1	0	1	123
	Grassland	58	3	64	694	72	4	0	895
	Agriculture or pasture	14	1	24	218	581	1	0	839
	Non-vegetated area	1	0	1	11	4	22	0	39
	River, lake or ocean	0	0	3	1	1	0	65	70
	Total	229	18	204	956	666	27	66	2166

Table 4. Contingency matrix and omission and commission errors resulting from comparing the 2001 map with the independent data set of the same year.

		Reference							Total
Classification	Classes	Natural woody vegetation	Forest plantation	Wetland	Grassland	Agriculture or pasture	Non-vegetated area	River, lake or ocean	
	Natural woody vegetation	146	14	3	26	8	0	0	197
	Forest plantation	2	19	0	4	1	0	0	26
	Wetland	8	1	91	9	8	0	7	124
	Grassland	28	8	42	569	170	1	1	819
	Agriculture or pasture	2	2	10	66	739	0	0	819
	Non-vegetated area	0	0	0	11	0	17	0	28
	River, lake or ocean	0	0	1	1	0	0	71	73
	Total	186	44	147	686	926	18	79	2086

Table 5. Contingency matrix and omission and commission errors resulting from comparing the 2018 map with the independent data set of the same year.

		Reference							Total
Classification	Classes	Natural woody vegetation	Forest plantation	Wetland	Grassland	Agriculture or pasture	Non-vegetated area	River, lake or ocean	
	Natural woody vegetation	150	9	2	21	6	0	0	188
	Forest plantation	3	55	0	1	4	0	0	63
	Wetland	7	1	85	6	9	0	7	115
	Grassland	24	9	39	438	193	1	1	705
	Agriculture or pasture	0	0	14	35	848	2	0	899
	Non-vegetated area	0	1	1	10	2	17	0	31
	River, lake or ocean	0	0	2	0	1	0	71	74
	Total	184	75	143	511	1063	20	79	2075




Figure 5. User and producer accuracies for each of mapped class in each evaluated year.

6 REFERENCES

- Baeza, S., Vélez-Martin, E., De Aballeyra, D., Banchemo, S., Gallego, F., Schirmbeck, J. & Hasenack, H. (2022). Two decades of land cover mapping in the Río de la Plata grassland region: The MapBiomass Pampa initiative. *Remote Sensing Applications: Society and Environment*, 28, 100834.
- Breiman, L. (2001). Random forests. *Machine learning*, v. 45, n. 1, p. 5-32.
- Liu F. T., Ting K. M., Zhou H. (2012). Isolation-based Anomaly Detection. *ACM Transactions on Knowledge Discovery from Data*, 6(1), 1556-4681.
- Olofsson, P., Foody, G. M., Herold, M., Stehman, S. V., Woodcock, C. E., & Wulder, M. A. (2014). Good practices for estimating area and assessing accuracy of land change. *Remote sensing of Environment*, 148, 42-57.
- Pontius Jr, R. G., & Millones, M. (2011). Death to Kappa: birth of quantity disagreement and allocation disagreement for accuracy assessment. *International Journal of Remote Sensing*, 32(15), 4407-4429.